

Influence the effect of long-range interaction on critical behavior of the three-dimensional systems.

S.V.Belim

Omsk State University

belim@univer.omsk.su

It is realized the theoretic - field description of Ising systems behaviour with effect of long-range interaction in two-loop approximation in three-dimensional space with using Padé-Borel resummation technique. The renorm-group equations are analysed and it is chosen fixed points, defining critical behaviour of the system. It is shown that the influence effect of long-range interaction can bring as to change the mode of the critical behaviour, so and to change the type of the phase transition.

Influence effect of long-range interaction, described on greater distances by sedate law $1/r^{-D-a}$ was explored analytically within the framework of ε -expansion approach [1, 2, 3] and numerically method Monte-Carlo [4, 5, 6] for one-dimensional and two-dimensional systems, and shown importance influences effect of long-range interaction on critical behaviour of Ising systems for value of the parameter $a < 2$. However before this times are absent article, in which given problem solved analytically at dimensionality space directly $D = 3$. However such description necessary on the strength of bad convergence of the rows, got within the framework of ε -expansion. In this article is conducted description of the critical behaviour three-dimensional Ising systems with effect of long-range interaction under different values of the parameter a .

Gamiltonian systems with provision for effect of long-range interaction:

$$H = \int d^D q \left\{ \frac{1}{2}(\tau_0 + q^a)\varphi^2 + u_0\varphi^4 \right\}, \quad (1)$$

here φ – fluctuations of the parameter of the order, D – dimension of space, $\tau_0 \sim |T - T_c|$, T_c – critical temperature, u_0 – positive constant. Critical behaviour greatly depends on parameter a , assigning velocity of the decrease the interaction with distance. As it is shown in work [1] influence effect of long-range interaction greatly under $0 < a < 2$, and for $a \geq 2$ critical behaviour equivalent of short-range interaction system. So hereinafter we limit event $0 < a < 2$.

Conducting standard renorm-groups procedure on base of the technology Feynmans diagrams [7] with $G(\vec{k}) = \frac{1}{\tau + |\vec{k}|^a}$, we get expressions for function β, γ_φ и γ_t , assigning the differential

renorm-group equation.

$$\begin{aligned}
\beta &= -(4-D)\left[1 - 36uJ_0 + 1728\left(2J_1 - J_0^2 - \frac{2}{9}G\right)u^2\right], \\
\gamma_t &= (4-D)\left[-12uJ_0 + 288\left(2J_1 - J_0^2 - \frac{1}{3}G\right)u^2\right], \\
\gamma_\varphi &= (4-D)96Gu^2, \\
J_1 &= \int \frac{d^D q d^D p}{(1 + |\vec{q}|^a)^2 (1 + |\vec{p}|^a) (1 + |q^2 + p^2 + 2\vec{p}\vec{q}|^{a/2})}, \\
J_0 &= \int \frac{d^D q}{(1 + |\vec{q}|^a)^2}, \\
G &= -\frac{\partial}{\partial |\vec{k}|^a} \int \frac{d^D q d^D p}{(1 + |q^2 + k^2 + 2\vec{k}\vec{q}|^a) (1 + |\vec{p}|^a) (1 + |q^2 + p^2 + 2\vec{p}\vec{q}|^{a/2})}
\end{aligned} \tag{2}$$

We shall redefy the efficient consn=tant of the interaction:

$$v = \frac{u}{J_0}. \tag{3}$$

As a result we come to following expression for function β , γ_φ and γ_t :

$$\begin{aligned}
\beta &= -(4-D)\left[1 - 36v + 1728\left(2\widetilde{J}_1 - 1 - \frac{2}{9}\widetilde{G}\right)v^2\right], \\
\gamma_t &= (4-D)\left[-12v + 288\left(2\widetilde{J}_1 - 1 - \frac{1}{3}\widetilde{G}\right)v^2\right], \\
\gamma_\varphi &= (4-D)96\widetilde{G}v^2, \\
\widetilde{J}_1 &= \frac{J_1}{J_0^2} \quad \widetilde{G} = \frac{G}{J_0^2}.
\end{aligned} \tag{4}$$

Such redefining have sense at values $a \leq D/2$. Herewith J_0 , J_1 , G become the dispersing function. Enterring parameter cutting Λ and considering limit of the relations

$$\begin{aligned}
\frac{J_1}{J_0^2} &= \frac{\int_0^\Lambda \int_0^\Lambda d^D q d^D p / ((1 + |\vec{q}|^a)^2 (1 + |\vec{p}|^a) (1 + |q^2 + p^2 + 2\vec{p}\vec{q}|^a))}{\left[\int_0^\Lambda d^D q / (1 + |\vec{q}|^a)^2\right]^2}, \\
\frac{G}{J_0^2} &= \frac{-\partial / (\partial |\vec{k}|^a) \int_0^\Lambda \int_0^\Lambda d^D q d^D p / ((1 + |q^2 + k^2 + 2\vec{k}\vec{q}|^a) (1 + |\vec{p}|^a) (1 + |q^2 + p^2 + 2\vec{p}\vec{q}|^a))}{\left[\int_0^\Lambda d^D q / (1 + |\vec{q}|^a)^2\right]^2}.
\end{aligned} \tag{5}$$

for $\Lambda \rightarrow \infty$ we get the finit expressions.

Values of integrals was found numerically. For $a \leq D/2$ sequence of values was built J_1/J_0^2 и G/J_0^2 for different values of Λ and was aproximated on infinity.

The mode of the critical behaviour is completely defined fixed points of renorm-group transformations, which can be found from condition equality zero β -function:

$$\beta(v^*) = 0. \tag{6}$$

The condition to stability is positive value of derived β -functions in fixed point:

$$\lambda = \frac{\partial\beta(v^*)}{\partial v} > 0. \quad (7)$$

Critical exponent ν , the characterizing growing of the radius of correlation in vicinities of the critical point ($R_c \sim |T - T_c|^{-\nu}$) is found as:

$$\nu = \frac{1}{2}(1 + \gamma_t)^{-1}.$$

Critical exponent η , the describing behaviour correlation functions in vicinities of the critical point in space wave vector ($G \sim k^{2+\eta}$), is defined on base scaling functions γ_φ : $\eta = \gamma_\varphi$. Over critical exponents can be determined from η and ν .

Known that expansion of theories of the indignations are asymptotic, but constants of the interaction of fluctuation parameter of the order is enough great to possible was directly use expressions (4).

So for the reason extractions from the got expressions to necessary physical information was an aplying method of the using Pade–Borel resummation technique. Herewith direct and reconversion Borels transformation are of the form:

$$f(v) = \sum_i c_i v^i = \int_0^\infty e^{-t} F(vt) dt, \quad (8)$$

$$F(v) = \sum_i \frac{c_i}{i!} v^i. \quad (9)$$

For calculations β -functions were used approximant Pade [2/1], for γ_t - and γ_φ -function – [1/1].

Firm fixed points renorm-groups transformations, derived β - functions in fixed to point and critical exponents for values of the parameter $1.5 \leq a \leq 1.9$ are provided in table. For values of the parameter $0 < a < 1.5$ exists only Gauss fixed point $v^* = 0$, not being firm. The givenned result will with predictions ε -expantion [1, 2, 3]. The absence firm fixed points is indicative of change the phase turning the second sort by phase turning the first type.

The analysis critical exponents shows that at reduction of the parameter a index ν decreases that is to say velocity of the growing of the radius correlation is slowed when approximation to critical point.

Thereby for three-dimensional Ising systems growing effect long-range interaction brings first to deceleration of the velocities of the growing of the radius корреляции in critical area, but then at importances of the parameter $a < 3/2$ to change the sort of the phase transition.

a	v^*	λ	ν	α	η	γ
1.5	0.015151	0.918690	0.555566	0.333302	0.002647	1.109661
1.6	0.015974	0.874129	0.557889	0.326333	0.003936	1.113582
1.7	0.020485	0.699732	0.567334	0.297998	0.004862	1.131910
1.8	0.023230	0.628209	0.572714	0.281858	0.007461	1.141155
1.9	0.042067	0.683927	0.620054	0.139838	0.013420	1.231787

Список литературы

- [1] *M. E. Fisher, S.-k. Ma and B. G. Nickel*, Phys. Rev. Lett. **29** , 917 (1972).
- [2] *J. Honkonen*, J. Phys. A **23** , 825 (1990).
- [3] *E. Luijten, H. Mebingfeld*, Phys. Rev. Lett. **86** , 5305 (2001).
- [4] *E. Bayong, H. T. Diep*. Phys. Rev. B. **9**, 18, 11920 (1999);
- [5] *E. Luijten*, Phys. Rev. E **60**, 7558 (1999).
- [6] *E. Luijten and H. W. J. Bloöte*, Phys. Rev. B **56**, 8945 (1997).
- [7] *J. Zinn-Justin*. Quantum field theory and critical phenomena, Clarendon Press, Oxford (1989).